

1. **Azaizeh, H.A., Neumann, G., and Marschner, H. (1995):** Effects of thiamine (vitamin B1) application on bulk soil and rhizosphere microorganisms and on thiamine released from developing bean (*Phaseolus vulgaris* L.) seedlings. *Z. Pflanzenernähr. Bodenk.* 158, 549-556.
2. **Azaizeh, H.A., Neumann, G., and Marschner, H. (1995):** Effects of thiamine on growth rates of *Rhizobium sp.* and diazotrophic bacteria cultivated in vitro at different pH. *Z. Pflanzenernähr. Bodenk.* 158, 557-562.
3. **Azaizeh, H.A., Neumann, G., and Marschner, H. (1996):** Effects of thiamine application and nitrogen fertilizer form on the number of N₂-fixing and total bacteria in the rhizosphere of maize plants. *Z. Pflanzenernähr. Bodenk.* 159, 183-188.
4. **Neumann, G., Azaizeh, H.A., and Marschner, H. (1996):** Thiamine (vitamin B1) seed treatment enhances germination and seedling growth of bean (*Phaseolus vulgaris* L.) exposed to soaking injury. *Z. Pflanzenernähr. Bodenk.* 159, 491-498.
5. **Neumann, G., Preißler, M., Azaizeh, H.A., and Römheld, V. (1999):** Thiamine (vitamin B1) deficiency in germinating seeds of *Phaseolus vulgaris* L. exposed to soaking injury. *Z. Pflanzenernähr. Bodenk.* 162, 295-300.
6. **Marschner, P., Neumann, G., Kania, A., Weiskopf, L., and Lieberei, R. (2002):** Spatial and temporal dynamics of the microbial community structure in the rhizosphere of cluster roots of white lupin (*Lupinus albus* L.) *Plant Soil* 246:167-174
7. **Wasaki, J., Rothe, A., Kania, A., Neumann, G., Römheld, V., Shinano, T., Osaki, M., and Kandeler, E. (2005):** Root exudation, P acquisition and microbial diversity in the rhizosphere of *Lupinus albus* as affected by P supply and atmospheric CO₂ concentration. *J. Environm. Qual.* 34, 2157-2166
8. **Weisskopf, L., Abou-Mansour, E., Fromin, N., Tomasi, N., Santelia, D., Edelkott, I., Neumann, G., Aragno, M., Tabacchi, R., and Martinoia, E. (2006):** White Lupin has developed a complex strategy to limit microbial degradation of secreted citrate required for phosphate nutrition. *Plant, Cell Environm.* 29, 919-927.
9. **Römheld, V., Neumann, G. (2006):** The Rhizosphere: Contributions of the soil-root interface to sustainable soil systems. In: Uphoff, N.; Ball, N.A.S.; Fernandes, E.; Herren, H.; Husson, O.; Laing, M.; Palm, C.; Thies, J. (eds), *Biological Approaches to Sustainable Soil Systems*, pp.92-107, CRC-Press, Oxford, UK
10. **Neumann, G. (2007):** Root exudates and nutrient cycling. In: *Soil Biology Vol. 10.* Marschner P., Rengel Z. (eds.) *Nutrient cycling in Ecosystems.* Springer, Berlin Heidelberg, pp.123-157.
11. **Haase, S., Neumann, G., Kania, A., Kuzyakov, Y., Römheld, V. and Kandeler, E. (2007):** Atmospheric CO₂ and the N-nutritional status modifies nodulation, nodule-carbon supply and root exudation of *Phaseolus vulgaris* L. *Soil Biol Biochem.* 39: 2208-2221.
12. **Haase, S., Ruess, L., Neumann, G., Marhan, S., Kandeler, E. (2008):** Low-level herbivory by root-rot nematodes (*Meloidogyne incognita*) modifies root hair morphology and rhizodeposition in host plants (*Hordeum vulgare*) *Plant Soil* 301: 151-164
13. **Römheld, V., Jiménez-Becker, S., Neumann, G., Gweyi-Onyango, J.P., Puelschen, L., Spreer, W., and Bangerth, F. (2008):** Non-nutritional fertigation effects as a challenge for improved production and quality in horticulture. In: *Fertigation: Optimizing the utilisation of water and nutrients.* International Potash Institute, Horgen, Switzerland., pp-103-115.

14. **Ögüt, M.; Er, F.;and Neumann, G. (2011):** Increased proton extrusion of wheat roots by inoculation with phosphorus solubilising microorganisms Plant and Soil, 339: 285-297.
15. **Cesco S, Mimmo T, Tonon G, Tomasi N, Pinton R, Terzano R, Neumann G, Weisskopf L, Renella G, Landi L, Nannipieri P (2012):** Plant-borne flavonoids released into the rhizosphere: impact on soil bio-activities related to plant nutrition. A review. Biol Fertil Soils 48: 123-149.
16. **Imran M, Kolla M, Römheld V, Neumann G (2013):** Impact of nutrient seed priming on germination, seedling development, nutritional status and grain yield of maize: J: Plant Nutr., in press.
17. **Imran M, Asim M, Römheld V, Neumann G (2013):** Nutrient seed priming improves seedling development and increases grain yield of maize exposed to low root zone temperatures during early growth. Europ. J. Agron.49: 141-148.
18. **Akter Z., Weinmann M., Neumann G., Römheld V. (2013):** An *in-vitro* .screening method to study the activity potential of biofertilizers based on Trichoderma and Bacillus sp. J. Plant Nutr. 36: 1439-1452.